



Golytely, Nulytely, Trilyte or Colyte (polyethylene glycol 3350 and electrolytes) Bowel Preparation Instructions

Bowel preparation before a colonoscopy actually starts days before, when you should stop taking certain medications or supplements. You must follow Golytely, Nulytely, Trilyte or Colyte instructions exactly for best results.

Key instructions

- Your bowel must be empty so that your doctor can clearly view your colon. Follow all of the instructions in this handout **EXACTLY** as they are written.
- DO NOT eat any solid food the ENTIRE day before your colonoscopy. Drink only clear liquids.
- Buy your bowel preparation at least **two days** before your colonoscopy.

Transportation on the day of your exam

- A responsible person **MUST BE PRESENT** with you at Check In before your colonoscopy and **MUST REMAIN** in the endoscopy area until you are discharged. You are not allowed to drive, take a taxi or bus, or leave the Endoscopy Center alone. **If you do not have a responsible driver (family member or friend) with you to take you home, your exam cannot be done with sedation and will be cancelled.**
- Please bring a list of all of your current medications, including any over-the-counter medications, with you.

Five (5) days before your colonoscopy

- Do **NOT** take medicines that stop diarrhea, such as loperamide (Imodium®) or bismuth subsalicylate (Kaopectate®, Pepto-Bismol®).
- Do **NOT** take fiber supplements, such as Metamucil®, Citrucel®, or Perdiem®.
- Do **NOT** take products that contain iron, such as multivitamins (the label lists what is in the products).

Buy the prescription bowel preparation solution at your local pharmacy or drugstore pharmacy two days before your colonoscopy.

Three days before your colonoscopy

- Do **NOT** eat high-fiber foods, such as popcorn, beans, seeds (flax, sunflower, quinoa), multigrain bread, nuts, salad/vegetables, or fresh and dried fruit.

One day before your colonoscopy

- **Only drink clear liquids the ENTIRE DAY before your colonoscopy. Do NOT eat any solid foods.** Drink at least **8 ounces** of clear liquids every hour after waking up. The chart shows what you can and cannot drink.

- **Clear Liquids
(No Red Liquids)**

DO NOT DRINK

Gatorade®, Pedialyte® or Powerade®	Alcohol
Clear broth or bouillon	Milk or non-dairy creamer
Coffee or tea (no milk or non-dairy creamer)	Noodles or vegetables in soup
Carbonated and non-carbonated soft drinks	Juice with pulp
Kool-Aid® or other fruit-flavored drinks	Liquid you cannot see through
Strained fruit juices (no pulp)	
Jell-O®, Popsicles®, hard candy	
Water	

The bowel preparation solution will be consumed in two parts.

Mix the solution the evening before your colonoscopy and refrigerate before drinking. You may add the flavor pack that came with the bowel preparation. Do NOT add ice, sugar or any other flavorings to the solution.

Part 1

At 6:00 pm the evening before your colonoscopy:

- Drink an 8-ounce glass of bowel preparation every 10 minutes for a total of 8 glasses.
- You may continue to drink clear liquids until midnight.

Part 2

SIX hours before your colonoscopy on the day of your procedure:

- Drink an 8-ounce glass of bowel preparation every 10 minutes for a total of 8 glasses.
- Fifteen minutes later, drink an 8-ounce glass of clear liquids every 15 minutes for a total of 2 glasses.
- You may continue to drink clear liquids up to four hours before your exam.